AA214 – Solution Set 2 Answer Sheet

1. Problem 1: Answers available upon request.

2. Problem 2

   (a)

$$\alpha\Delta x(\delta_x u)_j - (-u_{j+2} + \beta u_{j+1} - \beta u_{j-1} + u_{j-2}) = \alpha\Delta x\ er_t$$

	u_j	$\Delta x \left(\frac{\partial u}{\partial x}\right)_j$	$\Delta x^2 \left(\frac{\partial^2 u}{\partial x^2}\right)_j$	$\Delta x^3 \left(\frac{\partial^3 u}{\partial x^3}\right)_j$	$\Delta x^4 \left(\frac{\partial^4 u}{\partial x^4}\right)_j$	$\Delta x^5 \left(\frac{\partial^5 u}{\partial x^5}\right)_j$
$\alpha\Delta x(\delta_x u)_j$		α				
u_{j+2}	1	$\frac{(1)(2)^1}{1!}$	$\frac{(1)(2)^2}{2!}$	$\frac{(1)(2)^3}{3!}$	$\frac{(1)(2)^4}{4!}$	$\frac{(1)(2)^5}{5!}$
$-\beta u_{j+1}$	$-\beta$	$\frac{(-\beta)(1)^1}{1!}$	$\frac{(-\beta)(1)^2}{2!}$	$\frac{(-\beta)(1)^3}{3!}$	$\frac{(-\beta)(1)^4}{4!}$	$\frac{(-\beta)(1)^5}{5!}$
βu_{j-1}	β	$\frac{(\beta)(-1)^1}{1!}$	$\frac{(\beta)(-1)^2}{2!}$	$\frac{(\beta)(-1)^3}{3!}$	$\frac{(\beta)(-1)^4}{4!}$	$\frac{(\beta)(-1)^5}{5!}$
$-u_{j-2}$	-1	$\frac{(-1)(-2)^1}{1!}$	$\frac{(-1)(-2)^2}{2!}$	$\frac{(-1)(-2)^3}{3!}$	$\frac{(-1)(-2)^4}{4!}$	$\frac{(-1)(-2)^5}{5!}$
$\alpha\Delta x er_t$	0	0	0	0	0	?

   To get 4th irder accuracy, the first 5 columns must sum to 0. The $1^{st}, 3^{rd}$, and 5th columns are indentically 0, while setting the 2nd and 4th column to 0 give the two equations for the two unknowns α, β

$$\begin{aligned} \alpha + 4 - 2\beta &= 0 \\ 16 - 2\beta &= 0 \end{aligned}$$

   resulting in $\beta = 8$ and $\alpha = 12$ which gives the fourth order method with

$$er_t = \frac{1}{30}\Delta x^4 \left(\frac{\partial^5 u}{\partial x^5}\right)_j$$

   If $\beta = 4$, then $\alpha = 4$ and the fourth column does not sum to zero and the method is second order accurate.

3. From the Taylor table

	u_j	$\Delta x \cdot \left(\frac{\partial u}{\partial x}\right)_j$	$\Delta x^2 \cdot \left(\frac{\partial^2 u}{\partial x^2}\right)_j$	$\Delta x^3 \cdot \left(\frac{\partial^3 u}{\partial x^3}\right)_j$	$\Delta x^4 \cdot \left(\frac{\partial^4 u}{\partial x^4}\right)_j$
$a \cdot \Delta x \cdot \left(\frac{\partial u}{\partial x}\right)_{j-1}$		a	$a \cdot (-1) \cdot \frac{1}{1}$	$a \cdot (-1)^2 \cdot \frac{1}{2}$	$a \cdot (-1)^3 \cdot \frac{1}{6}$
$\Delta x \cdot \left(\frac{\partial u}{\partial x}\right)_j$		1			
$-d \cdot u_{j-1}$	$-d$	$-d \cdot (-1) \cdot \frac{1}{1}$	$-d \cdot (-1)^2 \cdot \frac{1}{2}$	$-d \cdot (-1)^3 \cdot \frac{1}{6}$	$-d \cdot (-1)^4 \cdot \frac{1}{24}$
$-c \cdot u_j$	$-c$				
$-b \cdot u_{j+1}$	$-b$	$-b \cdot (1) \cdot \frac{1}{1}$	$-b \cdot (1)^2 \cdot \frac{1}{2}$	$-b \cdot (1)^3 \cdot \frac{1}{6}$	$-b \cdot (1)^4 \cdot \frac{1}{24}$

the following equation has been constructed to maximize the order of accuracy

(a)

$$\begin{bmatrix} 0 & -1 & -1 & -1 \\ 1 & -1 & 0 & 1 \\ -2 & -1 & 0 & -1 \\ 3 & -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \\ 0 \\ 0 \end{bmatrix}$$

This has the solution $[a, b, c, d] = [2, 1, 4, -5]/4$, so the scheme can be expressed as

$$2(\delta_x u)_{j-1} + 4(\delta_x u)_j - \frac{1}{\Delta x}[-5u_{j-1} + 4u_j + u_{j+1}] = O(\Delta x^3)$$

The Taylor series error of this difference scheme is

$$\begin{aligned} er_t &= \left(2 \cdot (-1)^3 \cdot \frac{1}{6} + 5 \cdot (-1)^4 \cdot \frac{1}{24} - 1 \cdot (1)^4 \cdot \frac{1}{24}\right) \frac{\Delta x^3}{4} \left(\frac{\partial^4 u}{\partial x^4}\right)_j \\ &= -\frac{\Delta x^3}{24} \left(\frac{\partial^4 u}{\partial x^4}\right)_j \end{aligned}$$